

Your Tiopronin Tablets Savings Card

teva | tiopronin
tablets

Pay as
little as
\$0

Expires: **12/31/2026**

BIN **610020**

GRP **99994248**

PCN **N/A**

ID **63202507210**

You must present this offer and your primary insurance card to the pharmacist along with your prescription. Limitations apply. Please note this offer is not available for patients eligible for Medicare, Medicaid or any other public payer coverage. See full terms and conditions.

SAVINGS PROGRAM TERMS AND CONDITIONS

Terms, Conditions, and Eligibility Requirements: Eligible patients must have commercial prescription insurance with coverage for Teva's Tiopronin Tablets. Uninsured and cash-paying patients are NOT eligible for this Program. Patients enrolled in any state or federally funded healthcare program, including but not limited to, Medicare, Medigap, Medicaid, VA, DOD, TRICARE, Puerto Rico Government Health Insurance Plan, Medicare-eligible patients enrolled in an employer-sponsored health plan or prescription drug benefit program for retirees, are NOT eligible for this Program. Cash Discount Cards and other noninsurance plans are not valid as primary under this Program. This Program is restricted to residents of the United States and United States territories.

Patients may pay as little as \$0 out of pocket for Teva's Tiopronin Tablets. Maximum Program assistance per prescription and annual benefits apply and out-of-pocket expenses may vary. Patient is responsible for costs above maximum benefit amounts. This Program is not insurance. Void if copied, transferred, purchased, altered, or traded and where prohibited and restricted by law. The Program is not transferable. No substitutions are permitted. The Program form may not be sold, purchased, traded, or counterfeited. Void if reproduced. The Program benefit cannot be combined with any other financial assistance program, free trial, discount, prescription savings card, or other offer. This Program is managed by Mercialis on behalf of Teva Pharmaceuticals USA, Inc. Teva Pharmaceuticals USA, Inc. and its affiliates reserves the right to make eligibility determinations, to set Program benefit maximums, to monitor participation, and to change, rescind, revoke, or discontinue this Program at any time without notice. Limit one Program enrollment per individual. If you have any questions regarding this Program, your eligibility or benefits, or if you wish to discontinue your participation, please call 844-248-7949.

Expiration Date: 12/31/2026.

The Copay Card is intended for the benefit of patients, not their insurance plans or other third parties. Patients whose commercial insurance plans do not apply Copay Card payments to satisfy patient out-of-pocket cost sharing amounts may not be eligible for the Copay Card. Similarly, patients whose commercial insurance plans require use of the Copay Card as a condition of the plan waiving some or all of otherwise applicable patient out-of-pocket cost sharing amounts may not be eligible for the Copay Card or have a reduced annual maximum program benefit. If you believe your commercial insurance plan may have such limitations, please call 844-248-7949.

**Valid only for Teva's Tiopronin Tablets, National Drug Code:
00093-7909-01**

To the Patient: By redeeming this Program, you acknowledge that you are an Eligible Patient and you understand and agree to comply with the terms and conditions of this Program.

This Program is for eligible **Commercially Insured Patients only**. Patients may pay as little as \$0 out of pocket for Teva's Tiopronin Tablets. Maximum Program assistance per prescription and annual benefits apply and out-of-pocket expenses may vary. This Program must be presented along with your prescription for Tiopronin Tablets and your primary insurance card to participate in this Program. Program not valid for Non-Insured/Cash-Paying Patients or where Teva's Tiopronin Tablets are not covered by the primary insurance.

For patients residing in Maine and Maryland, please click [here](#).

To the Pharmacist: When you apply this Program, you are certifying that Teva's Tiopronin Tablets are being dispensed to an Eligible Patient in compliance with these terms and conditions and the Pharmacy has not submitted and will not submit a claim for reimbursement under any federal, state, or other governmental program for this prescription. For **Commercially Insured Patients**, please submit this claim to the primary Third-Party Payer first, then submit the balance due to Mercialis as a Secondary Payer COB (coordination of benefits) with patient responsibility and a valid Other Coverage Code (e.g., 08).

Reimbursement will be received from Mercialis. For questions regarding processing, please call the Help Desk at 844-248-7949.

Please see the accompanying full Prescribing Information on the following pages.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use TIOPRONIN TABLETS safely and effectively. See full prescribing information for TIOPRONIN TABLETS.

TIOPRONIN tablets, for oral use

Initial U.S. Approval: 1988

INDICATIONS AND USAGE

Tiopronin is a reducing and complexing thiol indicated, in combination with high fluid intake, alkali, and diet modification, for the prevention of cystine stone formation in adults and pediatric patients 9 years of age and older with severe homozygous cystinuria, who are not responsive to these measures alone. (1)

DOSAGE AND ADMINISTRATION

- The recommended initial dosage in adult patients is 800 mg/day. In clinical studies, the average dosage was about 1,000 mg/day. (2.1)
- The recommended initial dosage in pediatric patients 9 years of age and older is 15 mg/kg/day. Avoid dosages greater than 50 mg/kg per day in pediatric patients. (2.1, 5.1, 8.4)
- Administer tiopronin tablets in 3 divided doses at the same times each day at least one hour before or 2 hours after meals. (2.1)
- Measure urinary cystine 1 month after initiation of tiopronin tablets and every 3 months thereafter. (2.1)

DOSAGE FORMS AND STRENGTHS

Tablets: 100 mg (3)

CONTRAINDICATIONS

- Hypersensitivity to tiopronin or any component of tiopronin tablets. (4)

WARNINGS AND PRECAUTIONS

- Proteinuria, including nephrotic syndrome, and membranous nephropathy, has been reported with tiopronin use. Pediatric patients receiving greater than 50 mg/kg of tiopronin per day may be at increased risk for proteinuria. (2.1, 5.1, 8.4)
- Hypersensitivity Reactions have been reported during tiopronin treatment. (4, 5.2)

ADVERSE REACTIONS

Most common adverse reactions ($\geq 10\%$) are nausea, diarrhea or soft stools, oral ulcers, rash, fatigue, fever, arthralgia, proteinuria, and emesis. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Teva at 1-888-838-2872 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

USE IN SPECIFIC POPULATIONS

- Lactation: Breastfeeding is not recommended. (8.2)
- Geriatric: Choose dose carefully and monitor renal function in the elderly. (8.5)

Additional pediatric use information is approved for Mission Pharmacal Company's Thiola (tiopronin) tablets. However, due to Mission Pharmacal Company's marketing exclusivity rights, this drug product is not labeled with that information.

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 9/2024

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage

2.2 Monitoring

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

5.1 Proteinuria

5.2 Hypersensitivity Reactions

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

6.2 Postmarketing Experience

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Tiopronin tablets are indicated, in combination with high fluid intake, alkali, and diet modification, for the prevention of cystine stone formation in adults and pediatric patients 9 years of age and older with severe homozygous cystinuria, who are not responsive to these measures alone.

Additional pediatric use information is approved for Mission Pharmacal Company's Thiola (tiopronin) tablets. However, due to Mission Pharmacal Company's marketing exclusivity rights, this drug product is not labeled with that information.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage

Adults: The recommended initial dosage in adult patients is 800 mg/day. In clinical studies, the average dosage was about 1,000 mg/day.

Pediatrics: The recommended initial dosage in pediatric patients 9 years of age and older is 15 mg/kg/day. Avoid dosages greater than 50 mg/kg per day in pediatric patients [see Warnings and Precautions (5.1), Use in Specific Populations (8.4)].

Additional pediatric use information is approved for Mission Pharmacal Company's Thiola (tiopronin) tablets. However, due to Mission Pharmacal Company's marketing exclusivity rights, this drug product is not labeled with that information.

Administer tiopronin tablets in 3 divided doses at the same times each day at least one hour before or 2 hours after meals.

Consider starting tiopronin tablets at a lower dosage in patients with history of severe toxicity to d-penicillamine.

2.2 Monitoring

Measure urinary cystine 1 month after starting tiopronin tablets and every 3 months thereafter. Adjust tiopronin tablets dosage to maintain urinary cystine concentration less than 250 mg/L.

Assess for proteinuria before treatment and every 3 to 6 months during treatment [see Warnings and Precautions (5.1)].

Discontinue tiopronin tablets in patients who develop proteinuria, and monitor urinary protein and renal function. Consider restarting tiopronin tablets treatment at a lower dosage after resolution of proteinuria.

3 DOSAGE FORMS AND STRENGTHS

Tablets for oral use:

100 mg tablets: White to off-white round shaped, sugar coated tablets, imprinted with W on one side in black ink.

4 CONTRAINDICATIONS

Tiopronin is contraindicated in patients with hypersensitivity to tiopronin or any other components of tiopronin tablets [see Warnings and Precautions (5.2)].

5 WARNINGS AND PRECAUTIONS

5.1 Proteinuria

Proteinuria, including nephrotic syndrome, and membranous nephropathy, have been reported with tiopronin use. Pediatric patients receiving greater than 50 mg/kg of tiopronin per day may be at increased risk for proteinuria [see *Dosage and Administration (2.2)*, *Adverse Reactions (6.1, 6.2)*, *Use in Specific Populations (8.4)*]. Monitor patients for the development of proteinuria and discontinue therapy in patients who develop proteinuria [see *Dosage and Administration (2.2)*].

5.2 Hypersensitivity Reactions

Hypersensitivity reactions (drug fever, rash, fever, arthralgia and lymphadenopathy) have been reported [see *Contraindications (4)*].

6 ADVERSE REACTIONS

The following adverse reactions are discussed in greater detail in other sections of the labeling:

- Proteinuria [see *Warnings and Precautions (5.1)*]
- Hypersensitivity [see *Warnings and Precautions (5.2)*]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, the adverse reaction rates observed in the clinical trials of the drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Adverse reactions occurring at an incidence of $\geq 5\%$ in an uncontrolled trial in 66 patients with cystinuria age 9 to 68 years are shown in the table below. Patients in group 1 had previously been treated with d-penicillamine; those in group 2 had not. Of those patients who had stopped taking d-penicillamine due to toxicity (34 out of 49 patients in group 1), 22 were able to continue treatment with tiopronin. In those without prior history of d-penicillamine treatment, 6% developed reactions of sufficient severity to require tiopronin withdrawal.

Table 1 presents adverse reactions $\geq 5\%$ in either treatment group occurring in this trial.

Table 1: Adverse Reactions Occurring in One or More Patients

System Organ Class	Adverse Reaction	Group 1	Group 2
		Previously treated with d-penicillamine (N=49)	Naïve to d-penicillamine (N=17)
Blood and Lymphatic System Disorders	anemia	1 (2%)	1 (6%)
Gastrointestinal Disorders	nausea	12 (25%)	2 (12%)
	emesis	5 (10%)	–
	diarrhea/soft stools	9 (18%)	1 (6%)
	abdominal pain	–	1 (6%)
	oral ulcers	6 (12%)	3 (18%)
General Disorders and Administration Site Conditions	fever	4 (8%)	–
	weakness	2 (4%)	2 (12%)
	fatigue	7 (14%)	–
	peripheral (edema)	3 (6%)	1 (6%)
	chest pain	–	1 (6%)
Metabolism and Nutrition Disorders	anorexia	4 (8%)	–
Musculoskeletal and Connective Tissue Disorders	arthralgia	–	2 (12%)
Renal and Urinary Disorders	proteinuria	5 (10%)	1 (6%)
	impotence	–	1 (6%)
Respiratory, Thoracic and Mediastinal Disorders	cough	–	1 (6%)
Skin and Subcutaneous Tissue Disorders	rash	7 (14%)	2 (12%)

ecchymosis	3 (6%)	–
pruritus	2 (4%)	1 (6%)
urticaria	4 (8%)	–
skin wrinkling	3 (6%)	1 (6%)

Taste Disturbance

A reduction in taste perception may develop. It is believed to be the result of chelation of trace metals by tiopronin. Hypogeusia is often self-limited.

6.2 Postmarketing Experience

Adverse reactions have been reported from the literature, as well as during postapproval use of tiopronin. Because the postapproval reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to tiopronin exposure.

Adverse reactions reported during the postmarketing use of tiopronin are listed by body system in **Table 2**.

Table 2: Adverse Reactions Reported for Tiopronin Pharmacovigilance by System Organ Class and Preferred Term

System Organ Class	Preferred Term
Cardiac Disorders	congestive heart failure
Ear and Labyrinth Disorder	vertigo
Gastrointestinal Disorders	abdominal discomfort; abdominal distension; abdominal pain; chapped lips; diarrhea; dry mouth; dyspepsia; eructation; flatulence; gastrointestinal disorder; gastroesophageal reflux disease; nausea; vomiting; jaundice; liver transaminitis
General Disorders and Administration Site Conditions	asthenia; chest pain; fatigue; malaise; pain; peripheral swelling; pyrexia; swelling
Investigations	glomerular filtration rate decreased; weight increased
Metabolism and Nutrition Disorders	decreased appetite; dehydration; hypophagia
Musculoskeletal and Connective Tissue Disorders	arthralgia; back pain; flank pain; joint swelling; limb discomfort; musculoskeletal discomfort; myalgia; neck pain; pain in extremity
Nervous System Disorders	ageusia; burning sensation; dizziness; dysgeusia; headache; hypoesthesia
Renal and Urinary Disorders	nephrotic syndrome; proteinuria; renal failure
Skin and Subcutaneous Tissue Disorders	dry skin; hyperhidrosis; pemphigus foliaceus; pruritus; rash; rash pruritic; skin irritation; skin texture abnormal; skin wrinkling; urticaria

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Available published case report data with tiopronin have not identified a drug-associated risk for major birth defects, miscarriage, or adverse maternal or fetal outcomes. Renal stones in pregnancy may result in adverse pregnancy outcomes (*see Clinical Considerations*). In animal reproduction studies, there were no adverse developmental outcomes with oral administration of tiopronin to pregnant mice and rats during organogenesis at doses up to 2 times a 2 grams/day human dose (based on mg/m²). The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies are 2% to 4% and 15% to 20%,

respectively.

Clinical Considerations

Disease-associated maternal and/or embryo/fetal risk

Renal stones in pregnancy may increase the risk of adverse pregnancy outcomes, such as preterm birth and low birth weight.

Data

Animal Data

No findings of fetal malformations could be attributed to the drug in reproduction studies in mice and rats at doses up to 2 times the highest recommended human dose of 2 grams/day (based on mg/m²).

8.2 Lactation

Risk Summary

There are no data on the presence of tiopronin in either human or animal milk or on the effects of the breastfed child. A published study suggests that tiopronin may suppress milk production. Because of the potential for serious adverse reactions, including nephrotic syndrome, advise patients that breastfeeding is not recommended during treatment with tiopronin.

8.4 Pediatric Use

Tiopronin is indicated in pediatric patients 9 years of age and older with severe homozygous cystinuria, in combination with high fluid intake, alkali, and diet modification, for the prevention of cystine stone formation who are not responsive to these measures alone. This indication is based on safety and efficacy data from a trial in patients 9 years to 68 years of age and clinical experience. Proteinuria, including nephrotic syndrome, has been reported in pediatric patients. Pediatric patients receiving greater than 50 mg/kg tiopronin per day may be at greater risk [*see Dosage and Administration (2.1, 2.2), Warnings and Precautions (5.1), Adverse Reactions (6.1)*].

Tiopronin tablets are not approved for use in pediatric patients weighing less than 20 kg or in pediatric patients unable to swallow tablets [*see Dosage and Administration (2.1)*].

Additional pediatric use information is approved for Mission Pharmacal Company's Thiola (tiopronin) tablets. However, due to Mission Pharmacal Company's marketing exclusivity rights, this drug product is not labeled with that information.

8.5 Geriatric Use


This drug is known to be substantially excreted by the kidney, and the risk of adverse reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

10 OVERDOSAGE

There is no information on overdosage with tiopronin.

11 DESCRIPTION

Tiopronin immediate-release tablets are a reducing and cystine-binding thiol drug (CBTD) for oral use. Tiopronin is N-(2-Mercaptopropionyl) glycine and has the following structure:



Tiopronin has the empirical formula C₅H₉NO₃S and a molecular weight of 163.20. In this drug product tiopronin exists as a dl racemic mixture.

Tiopronin is a white to off-white crystalline powder, which is freely soluble in water.

Each tiopronin tablet contains 100 mg of tiopronin. The inactive ingredients in tiopronin tablets include colloidal silicon dioxide, corn starch, ethylcellulose, hydroxypropyl cellulose, lactose monohydrate, low substituted hydroxypropyl cellulose, magnesium stearate, silicified microcrystalline cellulose, and stearic acid. The tablets contain a coating that consists of glyceryl monocaprylocaprate, glyceryl monostearate, hypromellose 2910, medium chain triglycerides, polyvinyl alcohol-part hydrolyzed, polyethylene glycol 3350, sodium lauryl sulfate, sucrose, and talc. In addition, the imprinting ink contains ammonium hydroxide, black iron oxide, propylene glycol, and shellac.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The goal of therapy is to reduce urinary cystine concentration below its solubility limit. Tiopronin is an active reducing agent which undergoes thiol-disulfide exchange with cystine to form a mixed disulfide of tiopronin-cystine. From this reaction, a water-soluble mixed disulfide is formed and the amount of sparingly soluble cystine is reduced.

12.2 Pharmacodynamics

The decrement in urinary cystine produced by tiopronin is generally proportional to the dose. A reduction in urinary cystine of 250 to 350 mg/day at tiopronin dosage of 1 g/day, and a decline of approximately 500 mg/day at a dosage of 2 g/day, might be expected. Tiopronin has a rapid onset and offset of action, showing a fall in cystine excretion on the first day of administration and a rise on the first day of drug withdrawal.

12.3 Pharmacokinetics

Absorption

Tiopronin Tablets

When tiopronin single doses were given to fasted healthy subjects (n=39), the median time to peak plasma level (T_{max}) was 1 (range: 0.5 to 2.1) hours.

Elimination

Excretion

When tiopronin is given orally, up to 48% of dose appears in urine during the first 4 hours and up to 78% by 72 hours.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Long-term carcinogenicity studies in animals have not been performed.

Mutagenesis

Tiopronin was not genotoxic in the chromosomal aberration, sister chromatid exchange, and *in vivo* micronucleus assays.

Impairment of Fertility

High doses of tiopronin in experimental animals have been shown to interfere with maintenance of pregnancy and viability of the fetus. In 2 published male fertility studies in rats, tiopronin at 20 mg/kg/day intramuscular (IM) for 60 days induced reductions in testis, epididymis, vas deferens, and accessory sex glands weights and in the count and motility of cauda epididymal sperm.

16 HOW SUPPLIED/STORAGE AND HANDLING

100 mg: Each white to off-white round shaped, sugar coated tablet, imprinted with W on one side in black ink contains 100 mg of tiopronin. Tablets are available in bottles of 100 (NDC 0093-7909-01).

Store at 25°C (77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].

17 PATIENT COUNSELING INFORMATION

Lactation

Advise women that breastfeeding is not recommended during treatment with tiopronin tablets [*see Use in Specific Populations (8.2)*].

Manufactured In India By:

Watson Pharma Private Limited

Verna, Salcette Goa 403 722 India

Manufactured For:

Teva Pharmaceuticals

Parsippany, NJ 07054

Rev. A 9/2024